

Maximal residuated lattices with lifting Boolean center

George Georgescu¹, Laurențiu Leuştean^{2,3} and Claudia Mureşan¹

¹ Faculty of Mathematics and Computer Science, University of Bucharest,
Academiei 14, RO 010014, Bucharest, Romania
E-mails: ggeorgescu@rdslink.com, c.muresan@yahoo.com

² Department of Mathematics, Technische Universität Darmstadt,
Schlossgartenstrasse 7, 64289 Darmstadt, Germany

³ Institute of Mathematics "Simion Stoilow" of the Romanian Academy,
Calea Griviței 21, P.O. Box 1-462, Bucharest, Romania
E-mail: leustean@mathematik.tu-darmstadt.de

Abstract

In this paper we define, inspired by ring theory, the class of maximal residuated lattices with lifting Boolean center and prove a structure theorem for them: any maximal residuated lattice with lifting Boolean center is isomorphic to a finite direct product of local residuated lattices.

MSC: 06F35, 03G10.

Keywords: maximal residuated lattices, lifting Boolean center, semilocal residuated lattices, dense elements.

1 Introduction

A *commutative integral residuated bounded lattice* is an algebraic structure $(A, \vee, \wedge, \odot, \rightarrow, 0, 1)$ such that $(A, \vee, \wedge, 0, 1)$ is a bounded lattice, $(A, \odot, 1)$ is a commutative monoid and, for all $a, b, c \in A$,

$$a \leq b \rightarrow c \text{ if and only if } a \odot b \leq c.$$

Commutative integral residuated bounded lattices have been studied extensively and include important classes of algebras such as BL-algebras, introduced by Hájek as the algebraic counterpart of his Basic Logic [11], and MV-algebras, the algebraic setting for Łukasiewicz propositional logic (we refer to the monograph [5] for a detailed treatment of MV-algebras). Since in this paper we work only with commutative integral residuated bounded lattices, we shall call them simply *residuated lattices*. In order to simplify the notation, a residuated lattice

$(A, \vee, \wedge, \odot, \rightarrow, 0, 1)$ will be referred by its support set A . The *Boolean center* of A , denoted $B(A)$, is the set of all complemented elements of the bounded lattice $(A, \vee, \wedge, 0, 1)$.

The main purpose of this paper is to define the class of *maximal residuated lattices with lifting Boolean center* and to prove a structure theorem for them.

The inspiration for defining this class of residuated lattices comes from ring theory. Maximal rings are an important class of commutative rings with unit; we refer to [2] for a book treatment. The idea of *lifting idempotents*, due to Nicholson [16], turns out to be very useful in studying different classes of rings.

If A is a residuated lattice, $\{a_i\}_{i \in I} \subseteq A$ and $\{F_i\}_{i \in I}$ is a family of filters of A , then A is maximal iff, given a family of congruences $\{x \equiv a_i \pmod{F_i}\}_{i \in I}$ of A , being able to find a solution for any finite subset of these congruences implies one can find a solution for all the congruences. We refer to Section 6 for the formal definition.

Similar notions were developed for distributive lattices [10], MV-algebras [7] and BL-algebras [15]. Obviously, residuated lattices with a finite number of filters are maximal; hence, finite and simple residuated lattices are maximal. The converse is not true. An example of a maximal MV-algebra with an infinite number of ideals is given in [7, Proposition 9].

A residuated lattice A is said to have *lifting Boolean center* iff for every $e \in B(A/\text{Rad}(A))$ there exists a $f \in B(A)$ such that $e = f/\text{Rad}(A)$. Here $\text{Rad}(A)$ is the intersection of all maximal filters of A .

The main result of the paper is the following (see Theorem 6.6).

Theorem. Any maximal residuated lattice with lifting Boolean center is isomorphic to a finite direct product of local residuated lattices.

This structure theorem corresponds in the setting of residuated lattices to Zelinsky's theorem for maximal rings [17], [2, Theorem 2.6]. In fact, we prove even a stronger result, namely that a residuated lattice A with lifting Boolean center is maximal if and only if it is isomorphic to a finite direct product of some special residuated lattices, determined by elements in $B(A)$ (see Theorem 6.5).

2 Definitions and basic properties

We refer the reader to [9] for basic results in the theory of residuated lattices. In the following, we only present the material needed in the remainder of the paper.

We shall denote with $\mathcal{RL}$ the variety of residuated lattices and $\mathbf{RL}$ the category having as objects nontrivial residuated lattices and morphisms of residuated lattices as morphisms. We recall that by a *residuated lattice* we mean in fact a *commutative integral residuated bounded lattice*.

Let A be a residuated lattice. We use the notation $L(A)$ for the bounded

lattice $(A, \vee, \wedge, 0, 1)$. For all $a, b \in A$, let us define

$$\neg a := a \rightarrow 0, \quad a \leftrightarrow b := (a \rightarrow b) \wedge (b \rightarrow a).$$

The following lemma collects some useful properties (see for example [9]).

Lemma 2.1. *For all $a, b, c, d \in A$,*

- (i) $\neg 0 = 1, \neg 1 = 0$;
- (ii) $a = 1 \rightarrow a$ and $1 = a \rightarrow 1$;
- (iii) $a \leq b$ iff $a \rightarrow b = 1$ and $a \leq \neg b$ iff $a \odot b = 0$;
- (iv) if $a \leq b$ and $c \leq d$ then $a \odot c \leq b \odot d$;
- (v) if $a \leq b$ then $c \rightarrow a \leq c \rightarrow b$ and $b \rightarrow c \leq a \rightarrow c$;
- (vi) $a \odot b \leq a \wedge b$;
- (vii) $a \leq b \rightarrow a$;
- (viii) $a \odot 0 = 0, 0 \rightarrow a = 1$ and $a \leftrightarrow 0 = \neg a$;
- (ix) $a \leq \neg \neg a$ and $\neg \neg \neg a = \neg a$;
- (x) $a \odot (b \vee c) = (a \odot b) \vee (a \odot c)$;
- (xi) $(a \vee b) \rightarrow c = (a \rightarrow c) \wedge (b \rightarrow c)$;
- (xii) $a \rightarrow b \leq (c \rightarrow a) \rightarrow (c \rightarrow b)$;
- (xiii) $a \rightarrow (b \rightarrow c) = b \rightarrow (a \rightarrow c)$.

For all $a \in A$, we define $a^0 = 1$ and $a^n = a^{n-1} \odot a$ for all $n \in \mathbb{N}^*$. The *order* of $a \in A$, in symbols $ord(a)$, is the smallest $n \in \mathbb{N}$ such that $a^n = 0$. If no such n exists, then $ord(a) = \infty$. An element $a \in A$ is called: *nilpotent* iff $ord(a)$ is finite; a *unity* iff $\neg(a^n)$ is nilpotent for all $n \in \mathbb{N}$; *finite* iff both a and $\neg a$ are nilpotent.

A *filter* of A is a nonempty set $F \subseteq A$ such that, for all $a, b \in A$,

- (i) $a, b \in F$ implies $a \odot b \in F$;
- (ii) $a \in F$ and $a \leq b$ imply $b \in F$.

A filter F of A is *proper* iff $F \neq A$. We shall denote by $\mathcal{F}(A)$ the set of filters of A .

A proper filter P of A is called *prime* iff $a \vee b \in P$ implies $a \in P$ or $b \in P$ for all $a, b \in A$. The set of prime filters of A is denoted by $\text{Spec}(A)$.

A proper filter M of A is called *maximal* iff it is not contained in any other proper filter. We denote by $\text{Max}(A)$ the set of maximal filters of A . An immediate application of Zorn's lemma is the fact that any proper filter of A can be extended to a maximal filter. As a consequence, $\text{Max}(A) \neq \emptyset$ for any nontrivial residuated lattice A .

Let $X \subseteq A$. The filter of A generated by X will be denoted by $\langle X \rangle$. We have that $\langle \emptyset \rangle = \{1\}$ and for $X \neq \emptyset$,

$$< X > = \{a \in A \mid x_1 \odot \cdots \odot x_n \leq a \text{ for some } n \in \mathbb{N}^* \text{ and } x_1, \dots, x_n \in X\}.$$

For any $a \in A$, $< a >$ denotes the principal filter of A generated by $\{a\}$. Then $< a > = \{b \in A \mid a^n \leq b \text{ for some } n \in \mathbb{N}^*\}$.

The following results are standard and they are relatively easy to prove; for example, by following the proofs of the corresponding results for BL-algebras from [15, Chapter 1].

Proposition 2.2. $(\mathcal{F}(A), \subseteq)$ is a complete lattice. For every family $\{F_i\}_{i \in I}$ of filters of A , we have that

$$\bigwedge_{i \in I} F_i = \bigcap_{i \in I} F_i, \quad \bigvee_{i \in I} F_i = < \bigcup_{i \in I} F_i >.$$

The *radical* of A , denoted by $\text{Rad}(A)$, is the intersection of all maximal filters of A , when A is a nontrivial residuated lattice A . If $A = \{0\}$ is trivial, then $\text{Rad}(A) = \{0\}$ by definition.

Proposition 2.3. Let A be a residuated lattice. Then

- (i) $\text{Rad}(A) = \{a \in A \mid a \text{ is a unity}\};$
- (ii) $\text{Rad}(B) = B \cap \text{Rad}(A)$ for any subalgebra B of A ;
- (iii) $\text{Rad}(\prod_{i \in I} A_i) = \prod_{i \in I} \text{Rad}(A_i)$ for any family $\{A_i \mid i \in I\}$ of residuated lattices.

Proof. (i) See [12, Lemma 4.1]. (ii) and (iii) are easy consequences of (i). $\square$

A residuated lattice A is said to be *local* iff A has exactly one maximal filter. A local residuated lattice A is called *perfect* iff for all $a \in A$, $\text{ord}(a) < \infty$ if and only if $\text{ord}(\neg a) < \infty$.

Proposition 2.4. [6] A is local if and only if $D(A) := \{a \in A \mid \text{ord}(a) = \infty\}$ is the unique maximal filter of A .

The following lemma will be useful in Section 6.

Lemma 2.5. Define $F_M := \{a \in A \mid \text{the set } \{M \in \text{Max}(A) \mid a \notin M\} \text{ is finite}\}$. Then F_M is a filter of A and for any finite subset $\{M_1, \dots, M_n\}$ of $\text{Max}(A)$,

$$\bigcap \{M \mid M \in \text{Max}(A) - \{M_1, \dots, M_n\}\} \subseteq F_M.$$

Proof. We have that $1 \in F_M$, since $\{M \in \text{Max}(A) \mid 1 \notin M\} = \emptyset$. If $a, b \in A$, then

$$\{M \in \text{Max}(A) \mid a \odot b \notin M\} = \{M \in \text{Max}(A) \mid a \notin M\} \cup \{M \in \text{Max}(A) \mid b \notin M\},$$

hence $a, b \in F_M$ implies $a \odot b \in F_M$. If $a \leq b$, then

$$\{M \in \text{Max}(A) \mid b \notin M\} \subseteq \{M \in \text{Max}(A) \mid a \notin M\},$$

hence $a \in F_M$ implies $b \in F_M$.

If $a \notin F_M$, then the set $\{M \in \text{Max}(A) \mid a \notin M\}$ is infinite, hence there is $M \in \text{Max}(A) - \{M_1, \dots, M_n\}$ such that $a \notin M$. Thus, $a \notin \bigcap \{M \mid M \in \text{Max}(A) - \{M_1, \dots, M_n\}\}$. $\square$

Proposition 2.6. *Let B be a residuated lattice and A be a subalgebra of B . Then*

$$|\mathcal{F}(A)| \leq |\mathcal{F}(B)|, |\text{Spec}(A)| \leq |\text{Spec}(B)| \text{ and } |\text{Max}(A)| \leq |\text{Max}(B)|.$$

Proof. Just follow the proof of the corresponding result for BL-algebras [15, Proposition 1.2.25]. $\square$

If $h : A \rightarrow B$ is a morphism of residuated lattices, then the *kernel* of h is the set $\text{Ker}(h) := \{a \in A \mid h(a) = 1\}$. Obviously, h is injective iff $\text{Ker}(h) = \{1\}$.

Proposition 2.7. *Let $h : A \rightarrow B$ be a morphism of residuated lattices. Then the following properties hold:*

- (i) *for any (proper, prime, maximal) filter F of B , the set $h^{-1}(F) = \{a \in A \mid h(a) \in F\}$ is a (proper, prime, maximal) filter of A ; thus, in particular, $\text{Ker}(h)$ is a proper filter of A ;*
- (ii) *if h is surjective and F is a filter of A , then $h(F)$ is a filter of B ;*
- (iii) *if h is surjective and M is a maximal filter of A such that $h(M)$ is proper, then $h(M)$ is a maximal filter of B ;*
- (iv) *if h is surjective, then*

$$|\mathcal{F}(B)| \leq |\mathcal{F}(A)|, |\text{Spec}(B)| \leq |\text{Spec}(A)| \text{ and } |\text{Max}(B)| \leq |\text{Max}(A)|.$$

With any filter F of A we can associate a congruence relation $\equiv (\text{mod } F)$ on A by defining

$$a \equiv b (\text{mod } F) \text{ if and only if } a \leftrightarrow b \in F.$$

For any $a \in A$, let a/F be the equivalence class $a/\equiv (\text{mod } F)$. If we denote by A/F the quotient set $A/\equiv (\text{mod } F)$, then A/F becomes a residuated lattice with the operations induced from those of A .

Lemma 2.8. *Let F be a filter of A and $a, b \in A$. Then*

- (i) *$a/F = 1/F$ iff $a \in F$ and $a/F = 0/F$ iff $\neg a \in F$;*
- (ii) *$a/F \leq b/F$ iff $a \rightarrow b \in F$;*
- (iii) *if F is proper and $a/F = 0/F$, then $a \notin F$.*

The following proposition follows from a general result in universal algebra [4]. A proof for this particular case is similar to the proof of [7, Proposition 1].

Proposition 2.9. (Chinese Remainder Theorem)

Let $n \in \mathbb{N}^*$ and $F_1, \dots, F_n$ be filters of the residuated lattice A such that $F_i \vee F_j = A$ for every $i \neq j$, $i, j \in \overline{1, n}$. Then, for every $a_1, \dots, a_n \in A$, there exists an $a \in A$ such that $a \equiv a_i \pmod{F_i}$ for all $i \in \overline{1, n}$.

For any filter F of A , let us denote by p_F the quotient map from A onto A/F , defined by $p_F(a) = a/F$ for any $a \in A$. Then $F = \text{Ker}(p_F)$. For simplicity, we shall use the notation G/F for $p_F(G)$.

Lemma 2.10. Let F, G be filters of A such that $F \subseteq G$. Then

- (i) for all $a \in A$, $a/F \in G/F$ iff $a \in G$;
- (ii) G is proper iff G/F is a proper filter of A/F ;

Proposition 2.11. Let F, G be filters of A such that $F \subseteq G$. Then

- (i) the map p_F is an inclusion-preserving bijective correspondence between the filters of A containing F and the filters of A/F ; the inverse map is also inclusion-preserving;
- (ii) p_F maps the set of proper (maximal) filters of A containing F onto the set of proper (maximal) filters of A/F ;
- (iii) the map $\phi : A/F \rightarrow A/G$, $\phi(a/F) = a/G$ is a well-defined surjective morphism of residuated lattices; ϕ is an isomorphism if and only if $F = G$;
- (iv) the map $(A/F)/(G/F) \rightarrow A/G$, $(a/F)/(G/F) \mapsto a/G$ is an isomorphism of residuated lattices.

As an immediate application of Proposition 2.11.(ii) and Lemma 2.10, we get the following.

Proposition 2.12. Let A be a nontrivial residuated lattice and F be a proper filter of A .

- (i) Then $|\text{Max}(A/F)| \leq |\text{Max}(A)|$.
- (ii) Assume that $F \subseteq \text{Rad}(A)$. Then

$$|\text{Max}(A/F)| = |\text{Max}(A)| \text{ and } \text{Rad}(A/F) = \text{Rad}(A)/F.$$

In particular, A is local if and only if A/F is local.

Proposition 2.13. Let $\{A_i \mid i \in I\}$ be a family of residuated lattices and F_i be a filter of A_i for every $i \in I$. Then $F := \prod_{i \in I} F_i$ is a filter of $A := \prod_{i \in I} A_i$ and $A/F = \prod_{i \in I} A_i/F_i$. Moreover, if $p_F : A \rightarrow A/F$, $p_i : A_i \rightarrow A_i/F_i$ ($i \in I$) are the quotient maps, then $p_F = \prod_{i \in I} p_i$.

Let $B(A)$ be the Boolean center of A , that is the set of all complemented elements of the lattice $L(A)$. The following lemma collects some useful properties of $B(A)$.

Proposition 2.14. [3, 9]

- (i) $B(A)$ is a Boolean subalgebra of $L(A)$, $\neg e$ is the unique complement of $e \in B(A)$ and $\neg\neg e = e$.
- (ii) For any $e \in B(A)$, $e \odot e = e$ and $\langle e \rangle = \{a \in A \mid e \leq a\}$.
- (iii) For all $e, f \in B(A)$, $e \odot f = e \wedge f \in B(A)$, $e \rightarrow f = \neg e \vee f \in B(A)$ and $e \leftrightarrow f = (e \rightarrow f) \wedge (f \rightarrow e) \in B(A)$.
- (iv) $B(A) \cap \text{Rad}(A) = \{1\}$.
- (v) $B(\prod_{i \in I} A_i) = \prod_{i \in I} B(A_i)$ for any family $\{A_i \mid i \in I\}$ of residuated lattices.

Lemma 2.15. For every $e, f \in B(A)$ and $a, b \in A$, we have:

- (i) if $e \leq a$ then $\neg e \rightarrow a = a$;
- (ii) $e \rightarrow a = e \rightarrow (e \rightarrow a)$;
- (iii) $e \rightarrow (a \rightarrow b) = (e \rightarrow a) \rightarrow (e \rightarrow b)$;
- (iv) $\neg e \rightarrow a = e \vee a$;
- (v) $a \vee (e \wedge f) = (a \vee e) \wedge (a \vee f)$.

Proof. (i), (ii), (iii) and (iv) were proven by Kühr in [14] for bounded pseudo-BCK-algebras, hence for noncommutative residuated lattices. For the sake of completeness, we give the proofs in the commutative case.

- (i) $a = 1 \rightarrow a = (e \vee \neg e) \rightarrow a = (e \rightarrow a) \wedge (\neg e \rightarrow a) = \neg e \rightarrow a$, since $e \rightarrow a = 1$.

(ii)

$$\begin{aligned} e \rightarrow a &= 1 \rightarrow (e \rightarrow a) = (e \vee \neg e) \rightarrow (e \rightarrow a) \\ &= (e \rightarrow (e \rightarrow a)) \wedge (\neg e \rightarrow (e \rightarrow a)) = e \rightarrow (e \rightarrow a), \end{aligned}$$

since $\neg e = e \rightarrow 0 \leq e \rightarrow a$, so $\neg e \rightarrow (e \rightarrow a) = 1$.

- (iii) Since $a \leq e \rightarrow a$, we have that $a \rightarrow b \geq (e \rightarrow a) \rightarrow b$, hence

$$e \rightarrow (a \rightarrow b) \geq e \rightarrow ((e \rightarrow a) \rightarrow b) = (e \rightarrow a) \rightarrow (e \rightarrow b).$$

Furthermore, $a \rightarrow b \leq (e \rightarrow a) \rightarrow (e \rightarrow b)$ implies

$$\begin{aligned} e \rightarrow (a \rightarrow b) &\leq e \rightarrow ((e \rightarrow a) \rightarrow (e \rightarrow b)) = (e \rightarrow a) \rightarrow (e \rightarrow (e \rightarrow b)) \\ &= (e \rightarrow a) \rightarrow (e \rightarrow b), \text{ by (ii).} \end{aligned}$$

- (iv) Since $a \leq \neg e \rightarrow a$ and $e = \neg \neg e = \neg e \rightarrow 0 \leq \neg e \rightarrow a$, it follows that $e \vee a \leq \neg e \rightarrow a$.

Let $u \in A$ such that $u \geq e \vee a$. We get that

$$\begin{aligned} (\neg e \rightarrow a) \rightarrow u &\stackrel{(i)}{=} (\neg e \rightarrow a) \rightarrow (\neg e \rightarrow u) \stackrel{(iii)}{=} \neg e \rightarrow (a \rightarrow u) \\ &= \neg e \rightarrow 1 = 1. \end{aligned}$$

- (v) It is obvious that $a \vee (e \wedge f) \leq (a \vee e) \wedge (a \vee f)$. Now let $x \in A$ such that $x \leq (a \vee e) \wedge (a \vee f)$. Then, by (iv), $x \leq a \vee e = \neg e \rightarrow a$ and $x \leq a \vee f = \neg f \rightarrow a$, so, by the law of residuation, $x \odot \neg e \leq a$ and $x \odot \neg f \leq a$. It follows that

$$x \odot \neg(e \wedge f) = x \odot (\neg e \vee \neg f) = (x \odot \neg e) \vee (x \odot \neg f) \leq a,$$

hence, $x \leq \neg(e \wedge f) \rightarrow a = (e \wedge f) \vee a$, by (iv).

□

With the help of the Boolean center we can define a functor B between the category **RL** of residuated lattices and the category **Bool** of Boolean algebras as follows: for any morphism of residuated lattices $f : A_1 \rightarrow A_2$, $B(f) : B(A_1) \rightarrow B(A_2)$ is the restriction of f to $B(A_1)$.

For each $x \in A$, let us define the operation

$$\rightarrow_x : A \times A \rightarrow A, \quad a \rightarrow_x b = x \vee (a \rightarrow b).$$

Proposition 2.16. *Let F be a filter of A and $e \in B(A)$. Then*

- (i) $\langle e \rangle = \langle e, \vee, \wedge, \odot, \rightarrow_e, e, 1 \rangle$ is a residuated lattice;
- (ii) $F \cap \langle e \rangle = \{e \vee a \mid a \in F\}$ and $F \cap \langle e \rangle$ is a filter of $\langle e \rangle$;
- (iii) for all $a, b \in A$,

$$a \equiv b \pmod{F} \text{ in } A \quad \text{implies} \quad a \vee e \equiv b \vee e \pmod{F \cap \langle e \rangle} \text{ in } \langle e \rangle.$$

Proof. Just follow the proof for BL-algebras from [15, Proposition 1.4.4]. □

The following results are standard and they can be proved in a similar manner with the corresponding results for MV-algebras (see [5, Lemmas 6.4.4, 6.4.5]).

Proposition 2.17. *Let $\{A_i\}_{i \in I}$ be a nonempty family of residuated lattices and let $A \cong \prod_{i \in I} A_i$. Then there exists a family $\{e_i\}_{i \in I} \subseteq B(A)$ satisfying the following conditions:*

- (i) $\wedge_{i \in I} e_i = 0$;
- (ii) $e_i \vee e_j = 1$ whenever $i, j \in I, i \neq j$;
- (iii) each A_i is isomorphic to $\langle e_i \rangle$.

Proposition 2.18. *Let A be a residuated lattice, $n \geq 2$ and $e_1, \dots, e_n \in B(A)$ be such that*

- (i) $e_1 \wedge \dots \wedge e_n = 0$;
- (ii) $e_i \vee e_j = 1$ whenever $i, j \in \overline{1, n}, i \neq j$.

Then $A \cong \prod_{i=1}^n \langle \mathbf{e}_i \rangle$.

3 Finite direct products of residuated Lattices

In this section, we shall make a study of prime and maximal filters of finite direct products of residuated lattices, similar to the one done for MV-algebras in [1] or BL-algebras in [15].

In the sequel, I will be an index set, $\{A_i\}_{i \in I}$ a family of nontrivial residuated lattices and $A = \prod_{i \in I} A_i$. For each $i \in I$, let $pr_i : A \rightarrow A_i$, $pr_i((a_k)_{k \in I}) = a_i$ be the projections. Obviously, pr_i is a surjective morphism of residuated lattices. For any $i \in I$, let us denote by δ_i the element of A defined by

$$pr_i(\delta_i) = 0 \text{ and } pr_j(\delta_i) = 1 \text{ for all } j \in I, j \neq i.$$

It is clear that $\delta_i \vee \delta_k = 1$ for all $i, k \in I, i \neq k$.

Proposition 3.1. *Let P be a prime filter of A . Then*

- (i) $pr_i(P) \neq A_i$ for at most one $i \in I$;
- (ii) if I is finite, then $pr_i(P) \neq A_i$ for exactly one $i \in I$.

Proof.

- (i) Suppose that there are $i, k \in I, i \neq k$ such that $pr_i(P) \neq A_i$ and $pr_k(P) \neq A_k$. Then $\delta_i \vee \delta_k = 1 \in P$, so, $\delta_i \in P$ or $\delta_k \in P$, since P is a prime filter. It follows that $0 = pr_i(\delta_i) \in pr_i(P)$ or $0 = pr_k(\delta_k) \in pr_k(P)$. Since, by Proposition 2.7.(ii), $pr_i(P), pr_k(P)$ are filters, we get that $pr_i(P) = A_i$ or $pr_k(P) = A_k$. That is, we have got a contradiction.
- (ii) Let $I = \{1, \dots, n\}$. By (i), there exists at most one $i \in I$ such that $pr_i(P) \neq A_i$. Suppose that there is no such i , that is $pr_i(P) = A_i$ for all $i \in \overline{1, n}$. It follows that for every i there exists $a_i \in P$ such that $pr_i(a_i) = 0$. If we let $a := a_1 \odot \dots \odot a_n$, we get that $a \in P$ and $pr_i(a) = 0$ for all $i \in \overline{1, n}$, so $a = 0$. Thus, we have got that $0 \in P$, a contradiction.

□

If $i \in I$ and Q is a prime filter of A_i , $pr_i^{-1}(Q)$ is a prime filter of A , by Proposition 2.7.(i). We call it the *filter of A over Q* and we denote it by $Ov(Q)$. Let us define

$$\begin{aligned} Ov(A) &:= \{P \subseteq A \mid P = Ov(Q) \text{ for some } Q \in \bigcup_{i \in I} \text{Spec}(A_i)\} \\ &= \bigcup \{pr_i^{-1}(Q) \mid i \in I, Q \in \text{Spec}(A_i)\}. \end{aligned}$$

Then $Ov(A) \neq \emptyset$ and $Ov(A) \subseteq \text{Spec}(A)$.

Proposition 3.2. *Suppose that I is finite and P is a prime filter of A . Let $i \in I$ be unique such that $pr_i(P) \neq A_i$. Then*

- (i) *$pr_i(P)$ is a prime filter of A_i and, if P is maximal, then $pr_i(P)$ is also maximal;*
- (ii) *$P = Ov(pr_i(P))$.*

Proof. Let $I = \{1, \dots, n\}$.

- (i) By Proposition 2.7.(ii) and the fact that $pr_i(P) \neq A_i$, we get that $pr_i(P)$ is a proper filter of A_i . Let $x, y \in A_i$ be such that $x \vee y \in pr_i(P)$, so $x \vee y = pr_i(c)$ for some $c \in P$. Let us define $c_1, c_2 \in A$ by:

$$pr_i(c_1) = x, pr_i(c_2) = y \text{ and } pr_j(c_1) = pr_j(c_2) = pr_j(c) \text{ for all } j \neq i.$$

Since $c_1 \vee c_2 = c \in P$, we must have $c_1 \in P$ or $c_2 \in P$, hence $x \in pr_i(P)$ or $y \in pr_i(P)$.

If P is maximal, apply Proposition 2.7.(iii) to get that $pr_i(P)$ is maximal.

- (ii) $Ov(pr_i(P)) = pr_i^{-1}(pr_i(P)) \supseteq P$. It remains to prove the converse inclusion. Let $a \in Ov(pr_i(P))$. Since $pr_i(a) \in pr_i(P)$, $pr_i(q_i) = pr_i(a)$ for some $q_i \in P$. For $j \in I, j \neq i$, $pr_j(P) = A_j$, so there exists $q_j \in P$ such that $pr_j(q_j) = pr_j(a)$. Let

$$q := (\delta_1 \vee q_1) \odot \dots \odot (\delta_n \vee q_n).$$

Then $q \in P$ and $pr_j(q) = pr_j(q_j) = pr_j(a)$ for all $j \in I$, so $q = a$, hence $a \in P$.

□

We are now ready to prove the main result of this section.

Theorem 3.3. *Let $n \in \mathbb{N}^*$, $A_1, \dots, A_n$ be nontrivial residuated lattices and*

$$A = \prod_{i=1}^n A_i. \text{ Then}$$

$$\begin{aligned} \text{Spec}(A) &= \bigcup_{i=1}^n \{A_1 \times \dots \times A_{i-1} \times Q \times A_{i+1} \times \dots \times A_n \mid Q \in \text{Spec}(A_i)\}, \\ \text{Max}(A) &= \bigcup_{i=1}^n \{A_1 \times \dots \times A_{i-1} \times Q \times A_{i+1} \times \dots \times A_n \mid Q \in \text{Max}(A_i)\}, \end{aligned}$$

hence

$$|\text{Spec}(A)| = \sum_{i=1}^n |\text{Spec}(A_i)| \quad \text{and} \quad |\text{Max}(A)| = \sum_{i=1}^n |\text{Max}(A_i)|.$$

Proof. The first equality is an immediate application of Proposition 3.2; we get that $\text{Spec}(A) = \text{Ov}(A)$, hence the first equality.

If $M \in \text{Max}(A)$ and $i \in \overline{1, n}$ is unique such that $pr_i(M) \neq A_i$, then $pr_i(M) \in \text{Max}(A_i)$ by Proposition 3.2, and $M = \text{Ov}(pr_i(M)) = A_1 \times \dots \times A_{i-1} \times pr_i(M) \times A_{i+1} \times \dots \times A_n$. Conversely, if $Q \in \text{Max}(A_i)$ for some $i \in \overline{1, n}$, then $A_1 \times \dots \times A_{i-1} \times Q \times A_{i+1} \times \dots \times A_n = pr_i^{-1}(Q) \in \text{Max}(A)$ by Proposition 2.7.(i). $\square$

4 Dense elements and lifting Boolean center

Let A be a residuated lattice. An element a of A is said to be *dense* iff $\neg a = 0$. Following [8], we denote by $Ds(A)$ the set of the dense elements of A . It is easy to see that $Ds(A)$ is a filter of A satisfying $Ds(A) \subseteq \text{Rad}(A)$ [8].

Lemma 4.1. *Let A be a residuated lattice and $F \subseteq Ds(A)$ be a filter of A . Then*

- (i) *for all $a, b \in A$, $a/F = 0/F$ iff $a = 0$ and $a/F \leq \neg b/F$ iff $a \leq \neg b$;*
- (ii) *for all $a \in A$, $\text{ord}(a) = \text{ord}(a/F)$;*
- (iii) *$p_F(\{a \in A \mid \text{ord}(a) = \infty\}) = \{a/F \mid \text{ord}(a/F) = \infty\}$;*
- (iv) *for all $a \in A$, a is finite in A if and only if a/F is finite in A/F . Hence, $p_F(\{a \in A \mid a \text{ is finite}\}) = \{a/F \mid a/F \text{ is finite}\}$.*

Proof.

- (i) By Lemma 2.8.(i), $a/F = 0/F$ iff $\neg a \in F \subseteq Ds(A)$. Thus, $a/F = 0/F$ implies $\neg a \in Ds(A)$, so $\neg \neg a = 0$, that is equivalent to $a = 0$, since $a \leq \neg \neg a$. The converse implication is obvious.

$a/F \leq \neg b/F$ iff $a \leq \neg b$ follows using the above and the fact that in any residuated lattice $x \leq \neg y$ iff $x \odot y = 0$.

- (ii) By (i), for all $a \in A$ and all $n \in \mathbb{N}$, $a^n = 0$ iff $a^n/F = 0/F$ iff $(a/F)^n = 0/F$, hence $\text{ord}(A) = \text{ord}(a/F)$.

- (iii) $p_F(\{a \in A \mid \text{ord}(a) = \infty\}) = \{a/F \mid \text{ord}(a) = \infty\} \stackrel{(ii)}{=} \{a/F \mid \text{ord}(a/F) = \infty\}$.

- (iv) follows easily from (ii).

$\square$

Proposition 4.2. *Let A be a residuated lattice. Then*

(i) $Ds(A/F) = Ds(A)/F$ for any filter F of A contained in $Ds(A)$;

(ii) $Ds(B) = B \cap Ds(A)$ for any subalgebra B of A ;

(iii) $Ds\left(\prod_{i \in I} A_i\right) = \prod_{i \in I} Ds(A_i)$ for any family $\{A_i \mid i \in I\}$ of residuated lattices.

Proof. (i) is an immediate consequence of Lemma 4.1.(i). (ii), (iii) are obvious. $\square$

Proposition 4.3. *A residuated lattice A is local (perfect) if and only if $A/Ds(A)$ is local (perfect).*

Proof. Since $Ds(A) \subseteq \text{Rad}(A)$, we can apply Proposition 2.12 to get that A is local if and only if $A/Ds(A)$ is local. Use Proposition 4.1.(ii) to obtain the other equivalence. $\square$

Let $f : A \rightarrow B$ be a morphism of residuated lattices and define

$$\bar{f} : A/Ds(A) \rightarrow B/Ds(B), \quad \bar{f}(a/Ds(A)) = f(a)/Ds(B).$$

It is easy to see that $\bar{f}$ is a well-defined morphism of residuated lattices and that the diagram below is commutative (p_A and p_B are the canonical surjections).

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ p_A \downarrow & & \downarrow p_B \\ A/Ds(A) & \xrightarrow{\bar{f}} & B/Ds(B) \end{array}$$

As a consequence, we can define a (covariant) functor $\mathbf{T} : \mathbf{RL} \rightarrow \mathbf{RL}$ by setting $\mathbf{T}(A) = A/Ds(A)$ and $\mathbf{T}(f) = \bar{f}$.

Proposition 4.4. *$\mathbf{T}$ preserves surjective morphisms, injective morphisms and direct products.*

Proof. Let $f : A \rightarrow B$ be a morphism of residuated lattices. If f is surjective, then $p_B \circ f = \bar{f} \circ p_A$ is also surjective, hence $\bar{f}$ is surjective.

Assume now that f is injective and let $a_1, a_2 \in A$. Then we have the following sequence of equivalences: $\bar{f}(a_1/Ds(A)) = \bar{f}(a_2/Ds(A))$ iff $f(a_1)/Ds(B) = f(a_2)/Ds(B)$ iff $f(a_1) \leftrightarrow f(a_2) \in Ds(B)$ iff $f(a_1 \leftrightarrow a_2) \in Ds(B)$ iff $\neg f(a_1 \leftrightarrow a_2) = 0$ iff $f(\neg(a_1 \leftrightarrow a_2)) = 0$ iff (since f is injective) $\neg(a_1 \leftrightarrow a_2) = 0$ iff $a_1 \leftrightarrow a_2 \in Ds(A)$ iff $a_1/Ds(A) = a_2/Ds(A)$. Hence, $\bar{f}$ is injective.

Let $\{A_i \mid i \in I\}$ be a family of residuated lattices and $A = \prod_{i \in I} A_i$. By Proposition 4.2.(iii), $Ds(A) = \prod_{i \in I} Ds(A_i)$. Apply now Proposition 2.13. $\square$

A residuated lattice A is said to be *radical-dense* iff $\text{Rad}(A) = \text{Ds}(A)$. Let us denote with $\text{rd-}\mathcal{RL}$ the class of radical-dense residuated lattice. This terminology is inspired by [8], where a variety $\mathcal{A}$ is called *radical-dense* provided that A is a subvariety of $\mathcal{RL}$ and $\text{Rad}(A) = \text{Ds}(A)$.

Proposition 4.5. *rd-RL is closed with respect to subalgebras and direct products.*

Proof. Apply Propositions 4.2.(ii) and 2.3.(ii) to get closure under subalgebras. For obtaining closure with respect to direct products, use Propositions 4.2.(iii) and 2.3.(iii). $\square$

In the following, let A be a residuated lattice. Since $\text{Ds}(A) \subseteq \text{Rad}(A)$, we can apply Proposition 2.11.(iii) to get a surjective morphism of residuated lattices $\phi_A : A/\text{Ds}(A) \rightarrow A/\text{Rad}(A)$, $\phi_A(a/\text{Ds}(A)) = a/\text{Rad}(A)$ that makes the following diagram commutative.

$$\begin{array}{ccc} A & \xrightarrow{p_A} & A/\text{Ds}(A) \\ & \searrow r_A & \downarrow \phi_A \\ & & A/\text{Rad}(A) \end{array}$$

In the diagram above, p_A and r_A are the quotient maps. Moreover, ϕ_A is an isomorphism if and only if A is radical-dense, that is $\text{Ds}(A) = \text{Rad}(A)$.

If $B : \mathbf{RL} \rightarrow \mathbf{Bool}$ is the functor defined in Section 2, the diagram above induces the following commutative diagram in the category of Boolean algebras.

$$\begin{array}{ccc} B(A) & \xrightarrow{B(p_A)} & B(A/\text{Ds}(A)) \\ & \searrow B(r_A) & \downarrow B(\phi_A) \\ & & B(A/\text{Rad}(A)) \end{array}$$

Lemma 4.6. *$B(p_A)$ and $B(r_A)$ are injective.*

Proof. For all $e, f \in B(A)$, we have that $B(r_A)(e) = B(r_A)(f)$ iff $e/\text{Rad}(A) = f/\text{Rad}(A)$ iff $e \leftrightarrow f \in B(A) \cap \text{Rad}(A)$ iff $e \leftrightarrow f = 1$ (by Proposition 2.14.(iii), (iv)) iff $e = f$. Hence, $B(r_A)$ is injective and the fact that $B(p_A)$ is injective follows from the commutativity of the diagram. $\square$

We say that A has *lifting Boolean center* iff $B(r_A)$ is surjective (and hence a Boolean isomorphism).

Remark 4.7. *It is easy to see that the definition given above coincides with the one from the introduction: A has lifting Boolean center if and only if for every $e \in B(A/\text{Rad}(A))$ there exists an $f \in B(A)$ such that $e = f/\text{Rad}(A)$.*

The fact that MV-algebras have lifting Boolean center was already proved in [7, Proposition 5]. Moreover, using [15, Lemma 2.7.6], we can conclude that BL -algebras have lifting Boolean center too.

Proposition 4.8. *Any radical-dense residuated lattice has lifting Boolean center.*

Proof. If A is radical-dense, then ϕ_A is an isomorphism, by Proposition 2.11.(iii), hence $B(\phi_A)$ is an isomorphism of Boolean algebras. This implies obviously the surjectivity of $B(r_A)$. $\square$

Proposition 4.9. (i) *If A has lifting Boolean center then $B(\phi_A)$ is surjective.*

(ii) *$A/Ds(A)$ has lifting Boolean center if and only if $B(\phi_A)$ is a Boolean isomorphism.*

(iii) *Let $\{A_i \mid i \in I\}$ be a family of residuated lattices. Then $\prod_{i \in I} A_i$ has lifting Boolean center if and only if A_i has lifting Boolean center for every $i \in I$.*

Proof. (i) follows from the commutative diagram above.

(ii) For simplicity, we denote $A^* := A/Ds(A)$ and $H := \text{Rad}(A)/Ds(A)$. Since $Ds(A) \subseteq \text{Rad}(A)$, $\text{Rad}(A^*) = H$ by Proposition 2.12. Moreover, by applying Proposition 2.11.(iv) we obtain an isomorphism

$$\psi : A^*/H \rightarrow A/\text{Rad}(A), \quad \psi((a/Ds(A))/H) = a/\text{Rad}(A).$$


If $r_{A^*} : A^* \rightarrow A^*/\text{Rad}(A^*)$ is the quotient map, then $\psi \circ r_{A^*} = \phi_A$. By applying the functor B , we get that $B(\psi)$ is an isomorphism of Boolean algebras such that $B(\psi) \circ B(r_{A^*}) = B(\phi_A)$.

It follows that A^* has lifting Boolean center if and only if $B(r_{A^*})$ is an isomorphism if and only if $B(\phi_A)$ is an isomorphism.

(iii) Let $A := \prod_{i \in I} A_i$, $r_i : A_i \rightarrow A_i/\text{Rad}(A_i)$, $r_A : A \rightarrow A/\text{Rad}(A)$ be the quotient maps. Then $r_A = \prod_{i \in I} r_i$ by Propositions 2.13 and 2.3.(iii). Moreover, since $B(A) = \prod_{i \in I} B(A_i)$, it follows that $B(r_A) = \prod_{i \in I} B(r_i)$. $\square$

We finish this section with an example of a residuated lattice without lifting Boolean center.

Example 4.10. [9] Let $A = \{0, a, b, c, d, 1\}$ be the following residuated lattice.



$\rightarrow$	0	a	b	c	d	1
0	1	1	1	1	1	1
a	0	1	b	c	c	1
b	c	1	1	c	c	1
c	b	1	b	1	a	1
d	b	1	b	1	1	1
1	0	a	b	c	d	1

$\odot$	0	a	b	c	d	1
0	0	0	0	0	0	0
a	0	a	b	d	d	a
b	0	b	b	0	0	b
c	0	d	0	d	d	c
d	0	d	0	d	d	d
1	0	a	b	c	d	1

The maximal filters of A are $\{a, b, 1\}$ and $\{a, c, d, 1\}$, so $\text{Rad}(A) = \{a, 1\}$. It is easy to verify that $B(A) = \{0, 1\}$ and

$$B(A/\text{Rad}(A)) = A/\text{Rad}(A) = \{0/\text{Rad}(A), b/\text{Rad}(A), c/\text{Rad}(A), 1/\text{Rad}(A)\}.$$

Since $|B(A/\text{Rad}(A))| = 4 > 2 = |B(A)|$, $B(r_A)$ can not be surjective.

5 Semilocal residuated lattices

A residuated lattice is said to be *semilocal* iff it has only a finite number of maximal filters.

The trivial residuated lattice has no maximal filters, hence it is obviously semilocal. We shall consider only nontrivial semilocal residuated lattices. The class of semilocal residuated lattices includes finite residuated lattices as well as the local ones. It is easy to construct examples of semilocal residuated lattices that are not local: any finite direct product of $n \geq 2$ local residuated lattices has exactly n maximal filters, by Theorem 3.3.

Remark 5.1. *The class of semilocal residuated lattices is a pseudo-variety, i.e., it is closed under finite direct products, homomorphic images and subalgebras.*

Proof. Apply Theorem 3.3, Proposition 2.7.(iv) and Proposition 2.6. $\square$

Proposition 5.2. *Let A be a nontrivial residuated lattice and F be a proper filter such that $F \subseteq \text{Rad}(A)$. Then*

- (i) *A is semilocal if and only if A/F is semilocal;*
- (ii) *if A is a semilocal and $\text{Max}(A) = \{M_1, \dots, M_n\}$, then*

$$A/\text{Rad}(A) \cong \prod_{i=1}^n A/M_i.$$

Proof. (i) By Proposition 2.12.

(ii) Let us consider the canonical projection

$$\varphi : A/\text{Rad}(A) \rightarrow \prod_{i=1}^n A/M_i, \quad \varphi(a/\text{Rad}(A)) = (a/M_1, \dots, a/M_n).$$

For every $a, b \in A$, we get that $\varphi(a/\text{Rad}(A)) = \varphi(b/\text{Rad}(A))$ iff $a/M_i = b/M_i$ for all $i \in \overline{1, n}$ iff $a \leftrightarrow b \in M_i$ for all $i \in \overline{1, n}$ iff $a \leftrightarrow b \in \text{Rad}(A)$ iff $a/\text{Rad}(A) = b/\text{Rad}(A)$. Hence, φ is well defined and injective. Since $M_1, \dots, M_n$ are distinct maximal filters, it follows that $M_i \vee M_j = A$ for all $i \neq j$, so we can apply Proposition 2.9 to get for every $a_1, \dots, a_n \in A$ an $a \in A$ such that $a/M_i = a_i/M_i$ for all $i \in \overline{1, n}$, so $\varphi(a/\text{Rad}(A)) = (a_1/M_1, \dots, a_n/M_n)$. Thus, we have proved that φ is surjective too. It is easy to see that φ is a morphism of residuated lattices, hence it is an isomorphism of residuated lattices. $\square$

6 Maximal residuated lattices

Let A be a residuated lattice, I an index set, $\{a_i\}_{i \in I} \subseteq A$ and $\{F_i\}_{i \in I}$ be a family of filters of A . We say that the family $\{(a_i, F_i)\}_{i \in I}$ has *finite intersection property* (abbreviated *f.i.p.*) iff the family of sets $\{a_i/F_i\}_{i \in I}$ has finite intersection property, i.e. the intersection of every finite subfamily is nonempty. Formally, $\{(a_i, F_i)\}_{i \in I}$ has f.i.p. iff

for any finite $J \subseteq I$ there exists $x_J \in A$ with $x_J \equiv a_i(\text{mod } F_i)$ for all $i \in J$.

A is said to be *maximal* iff whenever $\{(a_i, F_i)\}_{i \in I}$ is a family with f.i.p., there exists $x \in A$ such that $x \equiv a_i(\text{mod } F_i)$ for all $i \in I$.

In this section we prove the main result of the paper, the structure theorem for maximal residuated lattices with lifting Boolean center. Before this, we give some useful properties of maximal residuated lattices.

By the Chinese Remainder Theorem, the following lemma is immediate.

Lemma 6.1. *Let A be a maximal residuated lattice. Then for any family $\{a_M\}_{M \in \text{Max}(A)}$ of elements of A there exists $x \in A$ such that $x \equiv a_M(\text{mod } M)$ for all $M \in \text{Max}(A)$.*

Proposition 6.2. *Any maximal residuated lattice is semilocal.*

Proof. Let A be a maximal residuated lattice, F_M be the filter of A defined in Lemma 2.5 and define $\mathcal{F} := \{(1, F_M)\} \cup \{(0, M) \mid M \in \text{Max}(A)\}$.

In order to prove that the family $\mathcal{F}$ has f.i.p., let us consider a finite subfamily $\{(1, F_M), (0, M_1), \dots, (0, M_n)\}$ and apply Lemma 6.1 to get the existence of an $x \in A$ satisfying $x \equiv 0(\text{mod } M_i)$ for all $i \in \overline{1, n}$ and $x \equiv 1(\text{mod } M)$ for all $M \in \text{Max}(A) - \{M_1, \dots, M_n\}$. By Lemma 2.8.(i),(iii) and Lemma 2.5, we get that $x \in \bigcap \{M \mid M \in \text{Max}(A) - \{M_1, \dots, M_n\}\} \subseteq F_M$, so $x \equiv 1(\text{mod } F_M)$. Since A is maximal and $\mathcal{F}$ has f.i.p., there exists $y \in A$ such that $y \equiv 1(\text{mod } F_M)$ and $y \equiv 0(\text{mod } M)$ for all $M \in \text{Max}(A)$. Thus, $y \in F_M$ and $y \notin M$ for any maximal filter M of A . It follows that $\text{Max}(A) = \{M \in \text{Max}(A) \mid y \notin M\}$, which is finite due to the fact that $y \in F_M$. $\square$

The converse of the above proposition does not hold: an example of a semilocal MV-algebra that is not maximal can be found in [7, Proposition 8].

Proposition 6.3. *Let A be a maximal residuated lattice and $e \in B(A)$. Then $\langle \mathbf{e} \rangle$ is also a maximal residuated lattice.*

Proof. Let $\mathcal{F} = \{(a_i, F_i)\}_{i \in I}$ be a family that has f.i.p. in $\langle \mathbf{e} \rangle$. Since $a_i \in A$ and every filter F_i of $\langle \mathbf{e} \rangle$ is also a filter of A , it follows that $\mathcal{F}$ has f.i.p. in A too. Apply now the fact that A is maximal to get an $x \in A$ such that $x \equiv a_i \pmod{F_i}$ in A for all $i \in I$. By Proposition 2.16.(iii), it follows that $x \vee e \equiv a_i \vee e \pmod{F_i \cap \langle e \rangle}$ in $\langle \mathbf{e} \rangle$ for all $i \in I$. Since $F_i \cap \langle e \rangle = F_i$ as $F_i \subseteq \langle e \rangle$ and $a_i \vee e = a_i$ as $a_i \in \langle e \rangle$, we get that $x \vee e \in \langle e \rangle$ is such that $x \vee e \equiv a_i \pmod{F_i}$ for all $i \in I$. Thus, $\langle \mathbf{e} \rangle$ is maximal. $\square$

Proposition 6.4. *The class of maximal residuated lattices is closed under finite direct products.*

Proof. Assume that $A = \prod_{i=1}^n A_i$, where $n \geq 1$ (the case $n = 0$ is trivial), where $A_1, \dots, A_n$ are maximal residuated lattices. By Proposition 2.17, there exist $e_1, \dots, e_n \in B(A)$ satisfying $e_1 \wedge \dots \wedge e_n = 0$, $e_i \vee e_j = 1$ for $i \neq j$ such that $A_i \cong \langle \mathbf{e}_i \rangle$ for all $i \in \overline{1, n}$. Let $\{(a_k, F_k)\}_{k \in K}$ have f.i.p. in A and apply Proposition 2.16.(iii) to get that the family $\{(a_k \vee e_i, F_k \cap \langle e_i \rangle)\}_{k \in K}$ has f.i.p. in $\langle \mathbf{e}_i \rangle$ for any $i \in \overline{1, n}$. Since $\langle \mathbf{e}_i \rangle$ is maximal, there exists $x_i \in \langle e_i \rangle$ such that $x_i \equiv a_k \vee e_i \pmod{F_k \cap \langle e_i \rangle}$ for all $k \in K$: in particular, $x_i \equiv a_k \vee e_i \pmod{F_k}$ in A for all $k \in K$. Let $x = x_1 \wedge \dots \wedge x_n$. Then $x \in A$ is such that $x \equiv (a_k \vee e_1) \wedge \dots \wedge (a_k \vee e_n) \pmod{F_k} \equiv a_k \pmod{F_k}$ for all $k \in K$, since, by Lemma 2.15.(v), $(a_k \vee e_1) \wedge \dots \wedge (a_k \vee e_n) = a_k \vee (e_1 \wedge \dots \wedge e_n) = a_k \vee 0 = a_k$. Thus, A is maximal. $\square$

Theorem 6.5. *Let A be a nontrivial residuated lattice with lifting Boolean center. Then the following are equivalent:*

(i) A is maximal;

(ii) there are $n \in \mathbb{N}^*$ and $e_1, \dots, e_n \in B(A)$ such that $A \cong \prod_{i=1}^n \langle \mathbf{e}_i \rangle$ and $\langle \mathbf{e}_i \rangle$ is a nontrivial maximal residuated lattice for all $i \in \overline{1, n}$.

Proof. (ii) $\Rightarrow$ (i) Apply Proposition 6.4.

(i) $\Rightarrow$ (ii) By Proposition 6.2, A is semilocal, hence $\text{Max}(A) = \{M_1, \dots, M_n\}$

for some $n \in \mathbb{N}^*$ and, moreover, $A/\text{Rad}(A) \cong \prod_{i=1}^n A/M_i$ (see Proposition 5.2).

Apply Proposition 2.17 to get $f_1, \dots, f_n \in B(A/\text{Rad}(A))$ such that $f_1 \wedge \dots \wedge f_n = 0/\text{Rad}(A)$, $f_i \vee f_j = 1/\text{Rad}(A)$ for $i \neq j$ and $A/M_i \cong \langle \mathbf{f}_i \rangle$ for all $i \in \overline{1, n}$. Since A has lifting Boolean center, there exist $e_1, \dots, e_n \in B(A)$ such that $f_i = e_i/\text{Rad}(A)$ for all $i \in \overline{1, n}$. It follows that $(e_1 \wedge \dots \wedge e_n)/\text{Rad}(A) = f_1 \wedge \dots \wedge f_n = 0/\text{Rad}(A)$, so $\neg(e_1 \wedge \dots \wedge e_n) \in B(A) \cap \text{Rad}(A) = \{1\}$, by Lemma 2.8.(i) and Proposition 2.14.(iv). Thus, $\neg(e_1 \wedge \dots \wedge e_n) = 1$, so $e_1 \wedge \dots \wedge e_n = \neg \neg(e_1 \wedge \dots \wedge e_n) = \neg 1 = 0$, by Proposition 2.14.(i). We get similarly that, for

$i \neq j$, $(e_i \vee e_j)/\text{Rad}(A) = f_i \vee f_j = 1/\text{Rad}(A)$, so $e_i \vee e_j = 1$. Applying now Proposition 2.18, it follows that $A \cong \prod_{i=1}^n \langle \mathbf{e}_i \rangle$. Moreover, $\langle \mathbf{e}_i \rangle$ is maximal for all $i \in \overline{1, n}$, by Proposition 6.3. Finally, $\langle \mathbf{e}_i \rangle$ is nontrivial, since A/M_i is nontrivial. $\square$

Theorem 6.6. *Let A be a nontrivial residuated lattice with lifting Boolean center. If A is maximal, then A is isomorphic to a finite direct product of local residuated lattices, each of which is clearly a maximal residuated lattice.*

Proof. By Theorem 6.5, $A \cong \prod_{i=1}^n \langle \mathbf{e}_i \rangle$, where $n = |\text{Max}(A)| \geq 1$, $e_1, \dots, e_n \in B(A)$ and $\langle \mathbf{e}_i \rangle$ is nontrivial and maximal for all $i \in \overline{1, n}$. It remains to prove that $\langle \mathbf{e}_i \rangle$ is local. Since $\langle \mathbf{e}_i \rangle$ is nontrivial, $|\text{Max}(\langle \mathbf{e}_i \rangle)| \geq 1$ for any $i \in \overline{1, n}$. On the other hand, applying Theorem 3.3, we get that $n = |\text{Max}(A)| = \sum_{i=1}^n |\text{Max}(\langle \mathbf{e}_i \rangle)| \geq n$. Thus, we must have $|\text{Max}(\langle \mathbf{e}_i \rangle)| = 1$ for all $i \in \overline{1, n}$, that is $\langle \mathbf{e}_i \rangle$ is local for all $i \in \overline{1, n}$. $\square$

References

- [1] L. P. Belluce, A. Di Nola, A. Lettieri, Subalgebras, direct products and associated lattices of MV-algebras, *Glasgow Math. J.* 34 (1992), 301-307.
- [2] W. Brandal, *Commutative rings whose finitely generated modules decompose*, Lecture Notes in Mathematics, 723, Springer-Verlag, Berlin, 1979.
- [3] D. Buşneag, D. Piciu, Residuated lattices of fractions relative to a $\wedge$ -closed system, *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)* 49 (97) (2006), 13-24.
- [4] S. Burris, H. P. Sankappanavar, *A course in universal algebra*, Graduate Texts in Mathematics, 78, Springer-Verlag, New York-Berlin, 1981.
- [5] R. Cignoli, I.M.L. D'Ottaviano, D. Mundici, *Algebraic foundations of many-valued reasoning*, Trends in Logic - Studia Logica Library, 7, Kluwer Acad. Publ., Dordrecht, 2000.
- [6] L. C. Ciungu, Classes of residuated lattices, *An. Univ. Craiova Ser. Mat. Inform.* 33 (2006), 189-207.
- [7] A. Filipoiu, G. Georgescu, A. Lettieri, Maximal MV-algebras, *Mathware Soft Comput.* 4 (1997), 53-62.
- [8] H. Freytes, Injectives in residuated algebras, *Algebra Universalis* 51 (2004), 373-393.

- [9] N. Galatos, P. Jipsen, T. Kowalski, H. Ono, *Residuated lattices: an algebraic glimpse at substructural logics*, Studies in Logic and The Foundations of Mathematics, 151, Elsevier, Amsterdam, 2007.
- [10] G. Georgescu, Finite products of local maximal lattices, *Math. Japonica* 37 (1991), 913-917.
- [11] P. Hájek, *Metamathematics of fuzzy logic*, Trends in Logic - Studia Logica Library, 4, Kluwer Acad. Publ., Dordrecht, 1998.
- [12] U. Höhle, Commutative residuated monoids, in: U. Höhle, P. Klement (eds.), *Non-classical logics and their applications to fuzzy subsets*, 53-106, Theory Decis. Lib. Ser. B Math. Statist. Methods, 32, Kluwer Acad. Publ., Dordrecht, 1995.
- [13] A. Iorgulescu, Classes of BCK Algebras-Part IV, Preprint Series of the Institute of Mathematics of the Romanian Academy 4 (2004), 1-37.
- [14] J. Kühr, Boolean and central elements and Cantor-Bernstein theorem in bounded Pseudo-BCK-algebras, accepted for publication in *J. Mult.-Valued Logic Soft Comput.*.
- [15] L. Leuştean, *Representations of many-valued algebras*, Ph. D. thesis, University of Bucharest, 2004.
- [16] W. K. Nicholson, Lifting Boolean center and exchange rings, *Trans. Amer. Math. Soc.* 229 (1977), 269-278.
- [17] D. Zelinski, Linearly compact modules and rings, *Amer. J. Math.* 75 (1953), 79-90.